

Graphs and Ideals generated by some 2-minors

MASAHIRO OHTANI

Graduate School of Mathematics, Nagoya University
Chikusa-ku, Nagoya 464-8602 JAPAN
m05011w@math.nagoya-u.ac.jp

Abstract

Let G be a finite graph on $[n] = \{1, 2, \dots, n\}$, X a $2 \times n$ matrix of indeterminates over a field K and $S = K[X]$ a polynomial ring over K . In this paper, we study about ideals I_G of S generated by 2-minors $[i, j]$ of X which correspond to edges $\{i, j\}$ of G . In particular, we construct a Gröbner basis of I_G as a set of paths of G and compute a primary decomposition.

1. Introduction

Let K be a field, $X = (X_{ij})$ an $m \times n$ generic matrix over K and $r \leq \min(m, n)$ a positive integer. The ideal $I_r(X)$ generated by *all* r -minors of X in a polynomial ring $S = K[X_{ij} \mid 1 \leq i \leq m, 1 \leq j \leq n]$ is called *the determinantal ideal* and is studied by many researchers from many different viewpoints. For example, $I_r(X)$ is a prime ideal in S and the quotient ring $S/I_r(X)$ is a Cohen-Macaulay ring, see Bruns and Herzog [1] or Bruns and Vetter [2].

In contrast, some kinds of ideals generated by *some* minors of X have been thought.

Conca defined ladder determinantal ideals in [3]. They are prime ideals and the quotient rings are Cohen-Macaulay.

Diaconis, Eisenbud and Sturmfels studied the ideal generated all “adjacent” 2-minors in a $2 \times n$ generic matrix [5]. An adjacent 2-minor of a $2 \times n$ matrix is the determinant of a submatrix with column indices j and $j + 1$ for $j = 1, 2, \dots, n - 1$. This ideal is generated by a regular sequence and is

not prime if $n > 2$. They compute a primary decomposition and all minimal prime ideals of them. Hosten and Sullivant studied *ideals of adjacent minors* as a generalization of ideals of Diaconis, Eisenbud and Sturmfels in [7]. An adjacent r -minor of X is the determinant of a submatrix with row indices $a_1, a_2, \dots, a_r$ and column indices $b_1, b_2, \dots, b_r$ where these indices are consecutive integers. They compute the minimal prime ideals of ideals of adjacent minors.

Diaconis, Eisenbud and Sturmfels also found a minimal primary decomposition of the ideal which is generated by all “corner minors” in [5]. A corner minor is the determinant of a 2×2 submatrix with row indices $1, i$ and column indices $1, j$.

In this paper, we study the following ideals generated by some 2-minors of a $2 \times n$ generic matrix.

From now on, we use the following notation. Let

$$X = \begin{pmatrix} X_1 & X_2 & \cdots & X_n \\ Y_1 & Y_2 & \cdots & Y_n \end{pmatrix}$$

be a $2 \times n$ generic matrix over a field K and $S = K[X_i, Y_i \mid 1 \leq i \leq n]$ a polynomial ring. For two integers $i, j \in [n] := \{1, 2, \dots, n\}$, we denote the 2-minor

$$\det \begin{pmatrix} X_i & X_j \\ Y_i & Y_j \end{pmatrix} = X_i Y_j - Y_i X_j$$

of X by $[i, j]$. Let G be a simple graph on $[n]$, i.e., G is a graph which does not have multiple edges or loops, and we define an ideal I_G as follows : ¹

$$I_G := ([i, j] \mid \{i, j\} \text{ is an edge of } G).$$

If G is a path, the ideal I_G coincides with the ideal of adjacent minors of X . If G is a star graph, then the ideal I_G coincides with the ideal of corner minors of X . So ideals I_G are generalizations of these ideals.

Properties of the ideal I_G are closely connected to properties of the graph G . The aim of this paper is to describe a Gröbner basis and a primary decomposition of I_G in relation to the datum of G . Particularly, we construct a Gröbner basis as “a set of paths” of G and describe an algorithm to compute a primary decomposition by “operations” of graphs.

¹After the first version of this paper was written, very recently, it has been brought to my attention that Herzog, Hibi, Hreinsdóttir, T. Kahle and J. Rauh wrote a paper [6] which has considerable overlaps with this paper.

This paper is organized as follows. Section 2 is preliminaries. We define some properties of paths, e.g., minimality and irreducibility, prove some facts and prepare some operations. In particular, decompositions of a path into a sum of irreducible paths play important roles in Section 3. In Section 3, we calculate a Gröbner basis of I_G as the set of all irreducible paths of G . In Section 4, we construct an “algorithm” of computation of a primary decomposition of I_G . In Section 5, we prove some results about connections between some properties of ideals I_G and graphs G . For example, we give a necessary condition for existence of Hamilton cycles of G .

ACKNOWLEDGMENT. The author is grateful to Mitsuyasu Hashimoto, Yuhi Sekiya and Ken-ichi Yoshida for valuable conversations and helpful suggestions. He also expresses his thanks to the referee for his many pieces of valuable advice. In particular, the example in Remark 5.6 is due to him.

The author was partially supported by JSPS Research Fellowships for Young Scientists.

2. Preliminaries

(2.1) A *walk* of G is a sequence $p_0e_0p_1e_1\ldots e_\ell p_\ell$ satisfying that each p_i is a vertex of G for $i = 0, 1, \ldots, \ell$, i.e., $p_i \in [n]$, that each e_i is an edge of G which connects vertices p_{i-1} and p_i for $i = 1, 2, \ldots, \ell$ and that $p_0 \leq p_\ell$. A walk $P = p_0e_0p_1e_1\ldots e_\ell p_\ell$ is a *path* if additionally it holds $p_i \neq p_j$ for each indices $i \neq j$. In this case, we call ℓ the *length* of P . A walk $P = p_0e_0p_1e_1\ldots e_\ell p_\ell e_{\ell+1}p_0$ is a *cycle* if the subsequence $p_0e_0p_1e_1\ldots e_\ell p_\ell$ is a path of G .

For a path $P = p_0e_1p_1e_2\ldots e_\ell p_\ell$, we call an element of $V(P) = \{p_i \mid 0 \leq i \leq \ell\}$ (resp. $J(P) = \{p_i \mid 0 < i < \ell\}$, and $E(P) = \{p_0, p_\ell\}$) a *vertex* (resp. a *joint*, and an *end*) of P .

(2.2) Now G is simple, so we can write $P = p_0p_1\ldots p_\ell$ short for a walk $P = p_0e_0p_1e_1\ldots e_\ell p_\ell$ with no confusion.

(2.3) Let $P = p_0p_1\ldots p_\ell$ be a walk. We take two vertices p_i and p_j with $i < j$. If $p_i < p_j$, we call the subsequence $p_i p_{i+1} \ldots p_j$ the *subwalk* of P from i to j , denoted $P_{i \rightarrow j}$. If $p_i > p_j$, the sequence $p_j p_{j-1} \ldots p_i$ is also called the subwalk of P from j to i , denoted $P_{j \rightarrow i}$.

(2.4) For two paths $P = p_0p_1\ldots p_\ell$ and $Q = q_0q_1\ldots q_m$ with $\#E(P) \cap E(Q) = 1$, we get the *sum* $P+Q$ of paths P and Q . For example, assume that $p_0 = q_0$ and $p_\ell < q_m$. Then the sum $P+Q$ is the walk $p_\ell p_{\ell-1} \ldots p_0 q_1 \ldots q_{m-1} q_m$.

(2.5) We define an order on the set of walks of G . For two walks $P = p_0 p_1 \dots p_\ell$ and $Q = q_0 q_1 \dots q_m$, we say that $P \leq Q$ if there is a sequence of indices $0 = j_0 < j_1 < \dots < j_\ell = m$ such that $q_{j_i} = p_i$ for each $i = 0, 1, \dots, \ell$.

A walk $P = p_0 p_1 \dots p_\ell$ is minimal with respect to this order if and only if P is a path and there never exists an edge which connects two vertices of P which do not adjoin.

(2.6) Let $P = p_0 p_1 \dots p_\ell$ be a minimal path of G . We say that P is *irreducible* if it holds that $J(P) \cap (p_0, p_\ell) = \emptyset$, where (p_0, p_ℓ) is the subset $\{x \mid p_0 < x < p_m\}$ of the set of real numbers $\mathbb{R}$.

(2.7) For an irreducible path $P = p_0 p_1 \dots p_\ell$, we set $g_P = M_P \cdot [p_0, p_\ell]$, where

$$M_P = \prod_{p \in J(P)} Z_p, \quad Z_p = \begin{cases} Y_p & \text{if } p < p_0 \\ X_p & \text{if } p > p_\ell \end{cases}.$$

(2.8) Let $P = p_0 p_1 \dots p_\ell$ be a minimal path of G . We can decompose P into a sum of irreducible paths.

We get a sequence $0 = i_0 < i_1 < \dots < i_s = \ell$ as follows. Set $i_0 := 0$ and

$$p_{i_t} := \begin{cases} \min J(P_{i_{t-1} \rightarrow \ell}) \cap (p_{i_{t-1}}, p_\ell) & \text{if } P_{i_{t-1} \rightarrow \ell} \text{ is not irreducible} \\ p_\ell & \text{if } P_{i_{t-1} \rightarrow \ell} \text{ is irreducible} \end{cases}$$

for $t > 0$. By definition, each subpath $P_{i_{t-1} \rightarrow i_t}$ is irreducible and $P = P_{i_0 \rightarrow i_1} + P_{i_1 \rightarrow i_2} + \dots + P_{i_{s-1} \rightarrow i_s}$ holds.

(2.9) Let $P = p_0 p_1 \dots p_\ell$ be an irreducible path and p_a a vertex of P . Assume that $p_a > p_m$. Then we decompose the subpath $P_{0 \rightarrow a}$ (or $P_{m \rightarrow a}$) into a sum $P_{i_0 \rightarrow i_1} + P_{i_1 \rightarrow i_2} + \dots + P_{i_{s-1} \rightarrow i_s}$ of irreducible paths, see (2.8).

Lemma 2.10. *Under the notation in (2.9), each monomial $M_{P_{i_{t-1} \rightarrow i_t}}$ for $t = 1, 2, \dots, s$, and each variable $X_{p_{i_t}}$ for $t = 1, 2, \dots, s-1$, divides M_P .*

Proof. It holds that $p_{i_t} > p_0$ by definition, so $p_{i_t} > p_m$ holds since P is irreducible. Then $X_{p_{i_t}}$ divides M_P .

Set $Q = P_{i_{t-1} \rightarrow i_t}$ and take a joint p_j of Q . It is enough to prove that M_P is divided by the variable Z_{p_j} if so is M_Q . If $p_j > p_{i_t} > p_m$, then the variable $Z_{p_j} = X_{p_j}$ divides M_Q and M_P . On the other hand, assume that $p_j < p_{i_{t-1}}$. By definition of $p_{i_{t-1}}$ (or earlier one), p_j must be less than p_0 . Then the variable $Z_{p_j} = Y_{p_j}$ divides M_Q and M_P . $\square$

(2.11) Let $P = p_0 p_1 \dots p_\ell$ and $Q = q_0 q_1 \dots q_m$ be irreducible paths of G with $p_0 = q_0$ and $p_\ell < q_m$. Take a minimal path $R = r_0 r_1 \dots r_k \leq P + Q$. We decompose R into a sum $R_{i_0 \rightarrow i_1} + R_{i_1 \rightarrow i_2} + \dots + R_{i_{s-1} \rightarrow i_s}$ of irreducible paths of G , see (2.8).

Lemma 2.12. *Under the notation in (2.11), each monomial $M_{R_{i_{t-1} \rightarrow i_t}}$ for $t = 1, 2, \dots, s$, and each variable $X_{r_{i_t}}$ for $t = 1, 2, \dots, s-1$, divides $\text{lcm}(M_P, M_Q) \cdot Y_{p_0}$.*

Proof. It holds that $r_{i_t} > p_\ell > p_0 = q_0$ by definition, so the variable $X_{r_{i_t}}$ divides M_P or M_Q .

We set $R' = R_{i_{t-1} \rightarrow i_t}$ and take a joint r_j of R' . It is enough to prove that $\text{lcm}(M_P, M_Q) \cdot Y_{p_0}$ is divided by the variable Z_{r_j} if so is $M_{R'}$. If $r_j > r_{i_t} > p_\ell$, then $r_j > p_0 = q_0$ and the variable $Z_{r_j} = X_{r_j}$ divides M_P or M_Q . On the other hand, assume that $r_j < r_{i_{t-1}}$. Then $r_j < r_{i_0} = p_\ell$ holds by definition of $r_{i_{t-1}}$ (or earlier one). So $r_j \leq p_0 = q_0$ holds and the variable $Z_{r_j} = Y_{r_j}$ divides M_P or M_Q or Y_{p_0} . $\square$

REMARK 2.13. We have analogues to discussions and results from (2.8) in the following situation.

Let $P = p_0 p_1 \dots p_\ell$ be a minimal path. We decompose P into a sum $P_{i_s \rightarrow i_{s-1}} + P_{i_{s-1} \rightarrow i_{s-2}} + \dots + P_{i_1 \rightarrow i_0}$ of irreducible paths of G as follows.

Set $i_0 = \ell$ and we inductively define indices i_t by

$$p_{i_t} := \begin{cases} \max J(P_{0 \rightarrow i_{t-1}}) \cap (p_0, p_{i_{t-1}}) & \text{if } P_{0 \rightarrow i_{t-1}} \text{ is not irreducible} \\ p_0 & \text{if } P_{0 \rightarrow i_{t-1}} \text{ is irreducible} \end{cases}.$$

(1) Let $P = p_0 p_1 \dots p_\ell$ be an irreducible path of G and p_a a joint of P with $p_a < p_0$. We decompose $P_{a \rightarrow 0}$ (or $P_{a \rightarrow m}$) into a sum $P_{i_s \rightarrow i_{s-1}} + P_{i_{s-1} \rightarrow i_{s-2}} + \dots + P_{i_1 \rightarrow i_0}$ of irreducible paths as above. Then each monomial $M_{P_{i_t \rightarrow i_{t-1}}}$ for $t = 1, 2, \dots, s$, and each variable $Y_{p_{i_t}}$ for $t = 1, 2, \dots, s-1$, divides M_P .

(2) Let $P = p_0 p_1 \dots p_\ell$ and $Q = q_0 q_1 \dots q_m$ be irreducible paths of G with $p_\ell = q_m$ and $p_0 < q_0$. Take a minimal path $R \leq P + Q$. We decompose R into a sum $R_{i_s \rightarrow i_{s-1}} + R_{i_{s-1} \rightarrow i_{s-2}} + \dots + R_{i_1 \rightarrow i_0}$ of irreducible paths as above. Then each monomial $M_{R_{i_t \rightarrow i_{t-1}}}$ for $t = 1, 2, \dots, s$, and each variable $Y_{r_{i_t}}$ for $t = 1, 2, \dots, s-1$, divides $\text{lcm}(M_P, M_Q) \cdot X_{p_\ell}$.


3. Gröbner Basis

(3.1) In this section, we use some definitions, properties and facts about Gröbner basis, e.g., monomial orders, definition of (reduced) Gröbner basis, and so on. We refer the reader to [4] for more information on them.

The following theorem is one of the main results of this paper.

Theorem 3.2. *The set $\mathcal{G} = \{g_P \mid P \text{ is an irreducible path of } G\}$ is the reduced Gröbner basis of I_G with respect to the reverse lexicographic order $<_{\text{revlex}}$ on S with $Y_1 > Y_2 > \dots > Y_n > X_1 > X_2 > \dots > X_n$.*

Example 3.3. Let G be the following graph :



A path of G is determined by its ends, so G has ten paths. But the Gröbner basis $\mathcal{G}$ of I_G consists of *nine* binomials since the path 2–4–1–5 is not irreducible. Explicitly,

$$\mathcal{G} = \left\{ \begin{array}{l} [1, 4], [1, 5], [2, 4], [3, 5], Y_1[4, 5], \\ X_4[1, 2], X_5[1, 3], Y_1X_5[3, 4], Y_1X_4X_5[2, 3] \end{array} \right\}$$

is the reduced Gröbner basis of I_G .

For the rest of this section, we prove Theorem 3.2.

(3.4) For an irreducible path $P = p_0p_1 \dots p_\ell$, the initial monomial $\text{in}_< g_P$ of g_P is $M_P \cdot X_{p_0}Y_{p_\ell}$.

Put $J(P) = \{p_{i_1} < p_{i_2} < \dots < p_{i_s} < p_{j_1} < p_{j_2} < \dots < p_{j_t}\}$ with $p_{i_s} < p_0 < p_\ell < p_{j_1}$. Then the binomial g_P is equal to

$$\prod_{1 \leq u \leq s} Y_{p_{i_u}} \cdot X_{p_0}Y_{p_m} \cdot \prod_{i \leq v \leq t} X_{p_{j_v}} - \prod_{1 \leq u \leq s} Y_{p_{i_u}} \cdot Y_{p_0}X_{p_m} \cdot \prod_{i \leq v \leq t} X_{p_{j_v}}.$$

Two terms of an element of $\mathcal{G}$ cannot be divided by the initial monomial of any element of $\mathcal{G}$, so $\mathcal{G}$ is reduced if $\mathcal{G}$ is a Gröbner basis of I_G .

(3.5) From now, we prove that $\mathcal{G}$ is a Gröbner basis of I_G by Buchberger's criterion.

In this and the next section, let R be a polynomial ring over a field K with a monomial order $<$. For two polynomials $f, g \in R$, then

$$S(f, g) = \frac{\text{in}_{<} g}{\gcd(\text{in}_{<} f, \text{in}_{<} g)} \cdot f - \frac{\text{in}_{<} f}{\gcd(\text{in}_{<} f, \text{in}_{<} g)} \cdot g$$

is called the S -polynomial of f and g .

Proposition 3.6 (Buchberger's criterion). *Let I be an ideal in R . A finite system of generators $\mathcal{G} = \{g_1, g_2, \dots, g_t\} \subset I$ is a Gröbner basis for I if and only if for all pairs $i \neq j$, the remainder on division of the S -polynomial $S(g_i, g_j)$ by $\mathcal{G}$ is zero.*

Proof. See Cox, Little and O'Shea [4], Chapter 2, §6, Theorem 6. $\square$

(3.7) First, we prove that $\mathcal{G} \subset I_G$.

Take an irreducible path $P = p_0 p_1 \dots p_\ell$. We prove it by induction on the length ℓ . If $\ell = 1$, P is an edge of G and $g_P = [p_0, p_\ell]$ is contained in I_G . Assume that $\ell > 1$. Then $J(P)$ is not empty, so either $J(P) \cap (-\infty, p_0)$ or $J(P) \cap (p_\ell, \infty)$ is not empty. If $J(P) \cap (-\infty, p_0) \neq \emptyset$ holds, we set $p_a := \max J(P) \cap (-\infty, p_0)$. Then subpaths $P_{a \rightarrow 0}$ and $P_{a \rightarrow \ell}$ are irreducible and $M_P = M_{P_{a \rightarrow 0}} \cdot M_{P_{a \rightarrow \ell}} \cdot Y_a$ holds. Then $g_P = Y_{p_0} M_{P_{a \rightarrow 0}} \cdot g_{P_{a \rightarrow \ell}} - Y_{p_\ell} M_{P_{a \rightarrow \ell}} \cdot g_{P_{a \rightarrow 0}}$ is contained in I_G . If $J(P) \cap (p_\ell, \infty) \neq \emptyset$, we can prove similarly for $a = \min J(P) \cap (p_\ell, \infty)$.

(3.8) We classify pairs of polynomials in $\mathcal{G}$ whose initial monomials are not coprime whether their ends are the same or not.

Let $P = p_0 p_1 \dots p_\ell$ and $Q = q_0 q_1 \dots q_m$ be irreducible paths of $\mathcal{G}$. Set $a = p_0$, $b = p_\ell$, $c = q_0$ and $d = q_m$. We suppose that $a \leq c$ without loss of generality. In addition, we assume that $\gcd(\text{in}_{<} g_P, \text{in}_{<} g_Q) \neq 1$.

(3.9) We first suppose that $a = c$ and $b = d$. Then the S -polynomial $S(g_P, g_Q)$ is equal to zero, so we have nothing to do.

(3.10) We secondly suppose that $a = c$ and $b \neq d$ hold. We can assume that $b < d$, then $S(g_P, g_Q) = \text{lcm}(M_P, M_Q) \cdot (Y_a X_b Y_d - Y_a Y_b X_d)$. We take a minimal path $R = r_0 r_1 \dots r_k \leq P + Q$ and decompose R into a sum $R_{i_0 \rightarrow i_1} + R_{i_1 \rightarrow i_2} + \dots + R_{i_{s-1} \rightarrow i_s}$ of irreducible paths as in (2.11). By Lemma 2.12,

$$S(g_P, g_Q) = M' \cdot X_b X_{r_{i_1}} X_{r_{i_2}} \dots X_{r_{i_{s-1}}} Y_d - Y_b X_{r_{i_1}} X_{r_{i_2}} \dots X_{r_{i_{s-1}}} X_d,$$

where M' is a monomial which can be divided by each $M_{R_{i_{t-1} \rightarrow i_t}}$ for $t = 1, 2, \dots, s$. Then it holds that

$$\begin{aligned} S(g_P, g_Q) &\stackrel{(s)}{\equiv} M' \cdot (X_b X_{r_{i_1}} X_{r_{i_2}} \cdots X_{r_{i_{s-2}}} Y_{r_{i_{s-1}}} X_d - Y_b X_{r_{i_1}} X_{r_{i_2}} \cdots X_{r_{i_{s-1}}} X_d) \\ &\stackrel{(s-1)}{\equiv} M' \cdot (X_b X_{r_{i_1}} X_{r_{i_2}} \cdots Y_{r_{i_{s-2}}} X_{r_{i_{s-1}}} X_d - Y_b X_{r_{i_1}} X_{r_{i_2}} \cdots X_{r_{i_{s-1}}} X_d) \\ &\equiv \cdots \equiv 0, \end{aligned}$$

where the equivalence (u) is induced by $g_{R_{i_{u-1} \rightarrow i_u}}$. As it were, the monomial $M_{R_{i_{t-1} \rightarrow i_t}}$ is a “catalyst” to exchange letters “ X ” and “ Y ”.

If $b = d$ and $a \neq c$, we can similarly prove using fact (2) in Remark 2.13.

(3.11) Until the end of this section, we suppose that $a \neq c$ and $b \neq d$. In this case, the following may hold only if $b < c$:

- (a) the variable Y_b divides M_Q , (b) the variable X_c divides M_P .

In fact, the condition (a) (resp. (b)) is equivalent to that $b \in J(Q)$ and $b < c$ (resp. $c \in J(P)$ and $b < c$).

(3.12) In this paragraph, we suppose that neither (a) nor (b) hold. Set $M'_P = \frac{M_P}{H}$ and $M'_Q = \frac{M_Q}{H}$, where $H = \gcd(M_P, M_Q)$. Then,

$$\begin{aligned} S(g_P, g_Q) &= H M'_P M'_Q (Y_a X_b X_c Y_d - X_a Y_b Y_c X_d) \\ &= M'_P Y_a X_b \cdot g_Q - M'_Q Y_c X_f \cdot g_P. \end{aligned}$$

(3.13) In this paragraph, we suppose that (a) holds and (b) does not hold. Now b is a joint of Q and $b < c$. So we decompose subpaths $Q_{b \rightarrow c}$ and $Q_{b \rightarrow d}$ into sums

$$\begin{aligned} Q_{b \rightarrow c} &= Q_{i_0 \rightarrow i_1} + Q_{i_1 \rightarrow i_2} + \cdots + Q_{i_{s-1} \rightarrow i_s}, \\ Q_{b \rightarrow d} &= Q_{j_0 \rightarrow j_1} + Q_{j_1 \rightarrow j_2} + \cdots + Q_{j_{t-1} \rightarrow j_t}. \end{aligned}$$

of irreducible paths of G as in (2.8). Set $M'_P = \frac{M_P}{H}$ and $M'_Q = \frac{M_Q}{Y_b H}$, where $H = \gcd(M_P, M_Q)$. Then,

$$\begin{aligned} S(g_P, g_Q) &= M'_P X_a \cdot g_Q - M'_Q X_c Y_d \cdot g_P \\ (\clubsuit) \quad &= H M'_P M'_Q (Y_a X_b X_c Y_d - X_a Y_b Y_c X_d). \end{aligned}$$

The first monomial of ($\clubsuit$) can be written as

$$N_1 \cdot \prod_{1 \leq v \leq t} M_{Q_{j_{v-1} \rightarrow j_v}} \cdot X_b Y_{j_1} Y_{j_2} \cdots Y_{j_{t-1}} Y_d,$$

where N_1 is a suitable monomial. Each $M_{Q_{j_{v-1} \rightarrow j_v}}$ is a “catalyst” to exchange letters “X” and “Y”, so it is equivalent to $H M'_P M'_Q \cdot Y_a Y_b X_c X_d$. Similarly, the second monomial of ($\clubsuit$) is equivalent to the same monomial by $g_{Q_{i_{u-1} \rightarrow i_u}}$ ’s.

Using fact (1) in Remark 2.13, $S(g_P, g_Q)$ is equivalent to 0 by $\mathcal{G}$ if (a) does not hold and (b) holds.

(3.14) In this paragraph, we suppose that (a) and (b) holds. Now b (resp. c) is a joint of Q (resp. P). Put $M'_P = \frac{M_P}{H X_c}$ and $M'_Q = \frac{M_Q}{H Y_b}$, where $H = \gcd(M_P, M_Q)$. Then $S(g_P, g_Q) = M'_Q Y_d g_P - M'_P X_a g_Q = H M'_P M'_Q (X_a Y_b Y_c X_d - Y_a X_b X_c Y_d)$. We decompose the subpath $P_{a \rightarrow c}$ (resp. $Q_{b \rightarrow d}$) into a sum $P_{i_0 \rightarrow i_1} + P_{i_1 \rightarrow i_2} + \cdots + P_{i_{s-1} \rightarrow i_s}$ (resp. $Q_{j_t \rightarrow j_{t-1}} + Q_{j_{t-1} \rightarrow j_{t-2}} + \cdots + Q_{j_1 \rightarrow j_0}$) of irreducible paths of P as in (2.9) (resp. in Remark 2.13 (1)). By Lemma 2.10 and the fact (1) in Remark 2.13, the monomials $M_{P_{i_{u-1} \rightarrow i_u}}$ and the variables $X_{p_{i_u}}$ (resp. $M_{Q_{j_v \rightarrow j_{v-1}}}$ and $Y_{j_{i_v}}$) divide M_P (resp. M_Q).

The S -polynomial $S(g_P, g_Q)$ is written as

$$(\spadesuit) \quad M_1 \cdot X_a X_{i_1} X_{i_2} \cdots X_{i_{s-1}} Y_c - M_2 \cdot X_b Y_{j_{t-1}} Y_{j_{t-2}} \cdots Y_{j_1} Y_d,$$

where M_1 (resp. M_2) is a monomial which is divided by $\prod_{1 \leq u \leq s} M_{P_{i_{u-1} \rightarrow i_u}}$ (resp. $\prod_{1 \leq v \leq t} M_{Q_{j_v \rightarrow j_{v-1}}}$). Each monomial in ($\spadesuit$) is equivalent to the monomial $H M'_P M'_Q Y_a Y_b X_c X_d$ by $g_{P_{i_{u-1} \rightarrow i_u}}$ ’s and $g_{Q_{j_v \rightarrow j_{v-1}}}$ ’s, so its remainder with respect to $\mathcal{G}$ is zero.

4. Primary Decomposition

Proposition 4.1. *The ideal I_G is a radical ideal.*

To prove Proposition 4.1, the following lemma is essential.

Lemma 4.2. *Let I be an ideal in a polynomial ring R over a field K . Assume that the initial ideal $\text{in}_< I$ with respect to a monomial order $<$ is generated by squarefree monomials. Then I is a radical ideal.*

Proof. Note that $\text{in}_< I$ is a radical ideal. Assume that $\sqrt{I} \neq I$ and take $f \in \sqrt{I} \setminus I$. Taking a normal form of f , we can suppose that $\text{in}_< f \notin \text{in}_< I$. Take an integer n with $f^n \in I$. Then $(\text{in}_< f)^n = \text{in}_< f^n \in \text{in}_< I$, so $\text{in}_< f \in \text{in}_< I$ and it is a contradiction. $\square$

By Proposition 4.1, the ideal I_G is the intersection of all prime ideals which contain I_G . In this section, we study a way to find all minimal prime ideals of I_G .

(4.3) Denote d_G the distance on the set of vertices of G which is defined by lengths of paths, namely, for two vertices x, y of G ,

$$d_G(x, y) := \min\{n \mid \text{there is a path } P \text{ of length } n \text{ with } E(P) = \{x, y\}\},$$

or $d_G(x, y) = \infty$ if there is not such a path.

(4.4) Let v be a vertex of G . We call the set $\{x \mid d_G(x, v) = 1\}$ the *neighborhood* of v in G , denoted $N_G(v)$. We say that G is *complete around* v if it holds that $d_G(x, y) \leq 1$ for any $x, y \in N_G(v)$.

(4.5) For a subset $A \subset [n]$, we denote the ideal $([a, b] \mid a, b \in A)$ in S by $I_2(A)$.

Proposition 4.6. *The following conditions are equivalent :*

- (1) G is complete around all vertices of G ,
- (2) G is a disjoint union of complete graphs,
- (3) the ideal I_G is a prime ideal.

Proof. (1) $\Rightarrow$ (2). Let x, y be vertices of G with $x < y$ and $d_G(x, y) < \infty$. It is enough to prove that $d_G(x, y) = 1$. Assume that $t := d_G(x, y) \geq 2$ and take a path $P = p_0 p_1 \dots p_t$ of length t with $p_0 = x$ and $p_t = y$. G is complete around p_1 , so there is an edge $\{p_0, p_2\}$ and it contradicts $d_G(x, y) = t$.

(2) $\Rightarrow$ (3) Put $G = \coprod_{1 \leq i \leq a} G_i$, where each G_i is a complete graph. We denote $V_i \subset [n]$ the vertex set of G_i . Then $I_G = \sum_i I_2(V_i)$ and

$$S/I_G \simeq \bigotimes_i \frac{k[X_v, Y_v \mid v \in V_i]}{I_2(V_i)}.$$

Each $k[X_v, Y_v \mid v \in V_i]/I_2(V_i)$ is a determinantal ring, so S/I_G is an integral domain.

(3) $\Rightarrow$ (1). Assume that there is a vertex v around which G is not complete. Then there are vertices $u, w \in N_G(v)$ with $d_G(u, w) = 2$. Then the equation

$$(\heartsuit) \quad 0 = \det \begin{pmatrix} X_u & X_v & X_w \\ X_u & X_v & X_w \\ Y_u & Y_v & Y_w \end{pmatrix} = X_u \cdot [v, w] - X_v \cdot [u, w] + X_w \cdot [u, v]$$

implies that $X_v \cdot [u, w] \in I_G$. I_G is not prime since $X_v \notin I_G$ and $[u, w] \notin I_G$. $\square$

(4.7) In this paragraph, we suppose that I_G is not prime. By Proposition 4.6, there is a vertex v around which G is not complete. Then there are $u, w \in N_G(v)$ with $d_G(u, w) = 2$.

Let P be a prime ideal of S which contains I_G . By the equation ($\heartsuit$), it holds that $X_v \in P$ or that $[u, w] \in P$. Similarly it holds that $Y_v \in P$ or that $[u, v] \in P$, so $(X_v, Y_v) \subset P$ or $[u, w] \in P$ holds. Thinking all pairs (u, w) of $N_G(v)$, P contains one of the following ideals :

$$(1) \quad I_G + (X_v, Y_v), \quad (2) \quad I_G + I_2(N_G(v)).$$

These ideals correspond to the following operations of graphs :

- ($\diamond$) (1) taking away v and all edges of which v is an end from G ,
(2) adding all edges which connect two vertices in $N_G(v)$ to G .

Lemma 4.8. *Let v be a vertex around which G is not complete, then*

$$I_G = (I_G + (X_v, Y_v)) \cap (I_G + I_2(N_G(v))).$$

Proof. The subsumption $\subset$ is trivial. Let P be a prime ideal which contains I_G . Then P contains the right hand side by the above discussion, so it holds that the right hand side is contained in $\sqrt{I_G} = I_G$. $\square$


Lemma 4.9. *In the above operations ($\diamond$), the number of vertices around which the graph is not complete decreases.*

Proof. Set G_1 (resp. G_2) as the graph which is made by the operation (1) (resp. (2)). Take a vertex $u \neq v$ of G . It is easy to see that G is not complete around u if so is G_1 .


Assume that G_2 is not complete around u . Then there are $x, y \in N_{G_2}(u)$ with $d_{G_2}(x, y) = 2$. We have nothing to prove if $d_G(x, y) = 2$. Otherwise, an edge $\{x, u\}$ or $\{y, u\}$ is added by the operation, i.e., it holds that $u \in N_G(v)$ and that x or $y \in N_G(v)$. If both x and $y \in N_G(v)$, then $d_{G_2}(x, y) = 1$. So $N_G(v)$ does not contain x or y . If $x \notin N_G(v)$, then x and $v \in N_G(u)$ and $d_G(x, v) = 2$. Then G is not complete around u . $\square$

By Lemma 4.8 and Lemma 4.9, repeating operations ($\diamond$), we have a set of prime ideals which contains all minimal prime ideals of I_G . So we can decompose I_G into the intersection of some prime ideals.


Example 4.10. Let G be the following graph :



Now, G is not complete around 1. Then we get the following graphs :



where a circled number means a vertex which is taken away. The left graph (1) is a disjoint union of complete graphs, but the right graph (2) is not complete around the vertex 2. So we operate on the graph (2) for $(\diamond)$, we get the following graphs.



Both graphs are disjoint unions of complete graphs. So the process finishes and we get the decomposition

$$I_G = (X_1, Y_1, [2, 3]) \cap (X_2, Y_2, [1, 4], [1, 5], [4, 5]) \cap I_2(X),$$

where $I_2(X)$ is the ideal generated by all 2-minors of X .

5. Properties of Graphs

In this section, we suppose that G is a connected simple graph on $[n]$ with $n > 2$.

Definition 5.1. A path P of G is *Hamilton* if $V(P) = [n]$. A cycle $p_0 p_1 \dots p_\ell p_0$ is *Hamilton* if the subpath $p_0 p_1 \dots p_\ell$ is a Hamilton path.

In graph theory, it is a very difficult and important problem whether G has a Hamilton cycle. In fact, a nontrivial necessary and sufficient condition for existence of Hamilton cycles is not known.

For example, G has a Hamilton cycle if one of the following holds :


- (1) $\sharp N_G(a) + \sharp N_G(b) \geq n$ for any vertices a and b , where $\sharp$ means the number of the elements of the set, see Ore [8].
- (2) G is planar and 4-connected, see Tutte [9].

Proposition 5.2. *If G has a Hamilton path, then $\text{ht } I_G = n - 1$. The converse holds if G is a tree, i.e., G has no cycle.*

Proof. Permuting vertices, we have a graph $G' \simeq G$ which has a Hamilton path $1-2-\dots-n$. Then I'_G contains a sequence $[1, 2], [2, 3], \dots, [n-1, n]$. It is a regular sequence and $\text{ht } I_G = \text{ht } I_{G'} \geq n - 1$. It holds that $\text{ht } I_G \leq n - 1$ since $I_G \subset I_2(X)$.

Assume that G is a tree. G has a vertex v with $\#N_G(v) \geq 3$ if G does not have a Hamilton path. Applying the operation (1) of $(\diamond)$ to v , we get a graph $\tilde{G}$ which has at most $n - 4$ edges. Then $\text{ht } I_{\tilde{G}} \leq n - 4$, so $\text{ht } I_G \leq \text{ht } I_{\tilde{G}} + 2 \leq n - 2$. $\square$

REMARK 5.3. If G is not a tree, the converse of Proposition 5.2 is not true in general. For example, let G be the following graph :




Then $\text{ht } I_G = 5$ but G does not have a Hamilton path.

(5.4) For an ideal I in S , we denote the set of prime ideals $\{P \in \text{Spec } S \mid P \supset I \text{ and } \text{ht } I = \text{ht } P\}$ by $\text{Assh}_S S/I$.

Proposition 5.5. *If G has a Hamilton cycle, then $\text{Assh}_S S/I_G = \{I_2(X)\}$.*

Proof. Take $P \in \text{Assh}_S S/I_G$. Then P is generated by some variables and some 2-minors of X , see (4.7). We suppose that P contains a variable X_v . Then $G \setminus v$ is a graph of size $n - 1$ and has a Hamilton path, so $\text{ht } I_{G \setminus v} = n - 2$. P contains $I_{G \setminus v} + (X_v, Y_v)$, then $\text{ht } P \geq \text{ht } I_{G \setminus v} + 2 = n$, it contradicts $\text{ht } P = \text{ht } I_G = n - 1$. $\square$

REMARK 5.6. The converse of Proposition 5.5 is not true in general. For example, let G be the following graph :



Now G has Hamilton paths, so $\text{ht } I_G = 6$ holds and $I_2(X)$ is contained in $\text{Assh}_S S/I_G$. However $G \setminus 7$ is isomorphic to the graph in Remark 5.3, and $G \setminus v$ has Hamilton paths for each $v = 1, 2, \dots, 6$. Then we have $\text{ht } I_{G \setminus v} = 5$ for each $v = 1, 2, \dots, 7$. So $\text{ht } P \geq 7$ holds for each associated prime ideal $P \neq I_2(X)$.

REFERENCES

- [1] W. Bruns and J. Herzog, Cohen-Macaulay rings, first paperback edition, Cambridge, 1998.
- [2] W. Bruns and U. Vetter, *Determinantal Rings, Lecture Notes in Mathematics*, **1327**, Springer-Verlag, 1988.
- [3] A. Conca, Ladder determinantal rings, *Journal of Pure and Appl. Algebra*, **98** (1995), 119–134.
- [4] D. Cox, J. Little and D. O’Shea, *Ideals, Varieties, and Algorithms*, third edition, Springer, 2006.
- [5] P. Diaconis, D. Eisenbud and P. Sturmfels, Lattice walks and primary decomposition, In Mathematical essays in honor of Gian-Carlo Rota (Cambridge, MA, 1996), pages 173–193. Birkhauser Boston, Boston, MA, 1998.
- [6] J. Herzog, T. Hibi, F. Hreinsdóttir, T. Kahle and J. Rauh, Binomial edge ideals and conditional independence statements, preprint, [arXiv:0909.4717](https://arxiv.org/abs/0909.4717)
- [7] S. Hosten and S. Sullivant, Ideals of Adjacent Minors, *J. Algebra*, **277** (2004), no.2., 615–642.
- [8] O. Ore, Note on Hamilton circuits, *Amer. Math. Monthly*, **67** (1960), no.1, 55.
- [9] W. T. Tutte, A theorem on planar graphs, *Trans. Amer. Math. Soc.*, **82** (1956), 99–116.